

## SOLVED EXERCISE 2.2

**1. Find the cube roots of -1, 8, -27, 64.**

**Solution:**

**(i) The three cube roots of -1**

$$\text{Let } x^3 = -1$$

$$(x)^3 + (1)^3 = 0$$

$$(x+1)(x^2 - x + 1) = 0$$

$$\text{Either } x+1=0 \quad \text{or} \quad x^2 - x + 1 = 0$$

$$x = -1$$

$$\text{Here } a = 1, b = -1, c = 1$$

Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{(-1) \pm \sqrt{(1)^2 - 4(1)(1)}}{2(1)}$$

$$x = \frac{1 \pm \sqrt{1-4}}{2}$$

$$x = \frac{1 \pm \sqrt{-3}}{2}$$

$$x = \frac{1 + \sqrt{-3}}{2} \quad \text{or} \quad x = \frac{1 - \sqrt{-3}}{2}$$

$$= -\left(\frac{-1 - \sqrt{-3}}{2}\right) \quad = -\left(\frac{-1 + \sqrt{-3}}{2}\right)$$

$$= -\omega^2 \quad = -\omega$$

Three cube roots of -1 are  $-1, -\omega, -\omega^2$

**(ii) The three cube roots of 8**

$$\text{Let } x^3 = 8$$

$$x^3 - 8 = 0$$

$$(x)^3 - (2)^3 = 0$$

$$(x-2)(x^2 + 2x + 4) = 0$$

$$\text{Either } x-2=0 \text{ or} \quad x^2 + 2x + 4 = 0$$

$$x = 2 \quad \text{or} \quad \text{Here } a = 1, b = 2, c = 4$$

Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-2 \pm \sqrt{(2)^2 - 4(1)(4)}}{2(1)}$$

$$x = \frac{-2\sqrt{4-16}}{2}$$

$$x = \frac{-2 \pm \sqrt{-12}}{2}$$

$$x = \frac{-2 \pm 2\sqrt{-3}}{2}$$

$$x = \frac{2(-1 \pm \sqrt{-3})}{2}$$

$$x = \frac{2(-1 \pm i\sqrt{3})}{2} \quad \because i = \sqrt{-1}$$

$$x = \frac{2(-1 + i\sqrt{3})}{2} \quad \text{or} \quad x = \frac{2(-1 - i\sqrt{3})}{2}$$

$$= 2\omega$$

$$= 2\omega^2$$

Three cube roots of 8 are  $2, 2\omega, 2\omega^2$

### (iii) The three cube roots of -27

$$\text{Let } x^3 = -27$$

$$x^3 + 27 = 0$$

$$(x)^3 - (3)^3 = 0$$

$$(x+3)(x^2 - 3x + 9) = 0$$

$$\text{Either } x+3 = 0 \quad \text{or} \quad x^2 - 3x + 9 = 0$$

$$x = -3$$

$$\text{Here } a = 1; b = -3, c = 9$$

Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-3 \pm \sqrt{(-3)^2 - 4(1)(9)}}{2(1)}$$

$$x = \frac{3 \pm \sqrt{9-36}}{2}$$

$$x = \frac{3 \pm \sqrt{-27}}{2}$$

$$x = \frac{3 \pm 3\sqrt{-3}}{2}$$

$$x = \frac{3(-1 \pm i\sqrt{3})}{2} \quad \because i = \sqrt{-1}$$

$$x = \frac{3(1 + i\sqrt{3})}{2} \quad \text{or} \quad x = 3\left(-\frac{1 - i\sqrt{3}}{2}\right)$$

$$x = \frac{-3(-1 - i\sqrt{3})}{2} \quad \text{or} \quad = -3\left(\frac{-1 + i\sqrt{3}}{2}\right)$$

$$= -3\omega^2 \quad = -3\omega$$

$\therefore$  Three cube roots of  $-27$  are  $-3, -3\omega, -3\omega^2$

#### (iv) The three cube roots of $64$

$$\text{Let } x^3 = 64$$

$$x^3 - 64 = 0$$

$$(x)^3 - (4)^3 = 0$$

$$(x - 4)(x^2 + 4x + 16) = 0$$

$$\text{Either } x - 3 = 0 \text{ or } x^2 + 4x + 16 = 0$$

$$x = 4$$

$$\text{Here } a = 1, b = 4, c = 16$$

Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-4 \pm \sqrt{(4)^2 - 4(1)(16)}}{2(1)}$$

$$x = \frac{-4 \pm \sqrt{16 - 64}}{2}$$

$$x = \frac{4 \pm \sqrt{-48}}{2}$$

$$x = \frac{-4 \pm 4\sqrt{-3}}{2}$$

$$x = \frac{4(-1 \pm i\sqrt{3})}{2} \quad i = \sqrt{-1}$$

$$x = \frac{3(-1 + i\sqrt{3})}{2} \quad \text{or} \quad x = 4\left(\frac{-1 - i\sqrt{3}}{2}\right)$$

$$= 4\omega \quad \quad \quad = 4\omega^2$$

Three cube roots of 64 are  $4, 4\omega, 4\omega^2$

## 2. Evaluate

(i)  $(1 - \omega - \omega^2)^7$

*Solution:*

$$\begin{aligned}(1 - \omega - \omega^2)^7 &= [1 - (\omega + \omega^2)]^7 \\&= [-1 - (-1)]^7 \quad \because \omega + \omega^2 = -1 \\&= (1 + 1)^7 \\&= 2^7 = 128\end{aligned}$$

(ii)  $(1 - 3\omega - 3\omega^2)^5$

*Solution:*

$$\begin{aligned}(1 - 3\omega - 3\omega^2)^5 &= [1 - 3(\omega + \omega^2)]^5 \\&= [1 - 3(-1)]^5 \quad \because \omega + \omega^2 = -1 \\&= (1 + 3)^5 \\&= 4^5 = 1024\end{aligned}$$

(iii)  $(9 + 4\omega + 4\omega^2)^3$

*Solution:*

$$\begin{aligned}(9 + 4\omega + 4\omega^2)^3 &= [9 + 4(\omega + \omega^2)]^3 \\&= [9 + 4(-1)]^3 \quad \because \omega + \omega^2 = -1 \\&= (9 - 4)^3 \\&= 5^3 = 125\end{aligned}$$

(iv)  $(2 + 2\omega - 2\omega^2)(3 - 3\omega + 3\omega^2)$

*Solution:*

$$\begin{aligned}(2 + 2\omega - 2\omega^2)(3 - 3\omega + 3\omega^2) \\&= [2(1 + \omega) - 2\omega^2][3(1 + \omega^2) - 3\omega] \\&\doteq [2(-\omega^2) - 2\omega^2][3(-\omega) - 3\omega] \quad \because 1 + \omega = -\omega^2\end{aligned}$$

$$\begin{aligned}
&= (-2\omega^2 - 2\omega^2)(-3\omega - 3\omega) && \because 1 + \omega^2 = -\omega \\
&= (-4\omega^2)(-6\omega) \\
&= (-4)(-6)(\omega^2 \cdot \omega) \\
&= 24\omega^3 \\
&= 24(1) && \because \omega^3 = 1 \\
&= 24
\end{aligned}$$

(v)  $(-1 + \sqrt{-3})^6 + (-1 - \sqrt{-3})^6$

*Solution:*

$$\begin{aligned}
&(-1 + \sqrt{-3})^6 + (-1 - \sqrt{-3})^6 \\
&= (2\omega)^6 + (2\omega^2)^6 && \because \omega = \frac{-1 + \sqrt{-3}}{2} \quad \text{and} \quad \omega^2 = \frac{-1 - \sqrt{-3}}{2} \\
&= 2^6 (\omega^6) + 2^6 (\omega^{12}) && 2\omega = -1 + \sqrt{-3} \quad 2\omega^2 = -1 - \sqrt{-3} \\
&= 2^6 [(\omega^3)^2] + 2^6 [(\omega^3)^4] \\
&= 2^6 [(1)^2 + (1)^4] && \because \omega^3 = 1 \\
&= 2^6 [1 + 1] \\
&= 2^6 \cdot 2 = 2^{6+1} = 2^7 \\
&= 128
\end{aligned}$$

(vi)  $\left(\frac{-1 + \sqrt{-3}}{2}\right)^9 + \left(\frac{-1 - \sqrt{-3}}{2}\right)^9$

*Solution:*

$$\begin{aligned}
&\left(\frac{-1 + \sqrt{-3}}{2}\right)^9 + \left(\frac{-1 - \sqrt{-3}}{2}\right)^9 \\
&= \omega^9 + (2\omega^2)^9 && \because \omega = \frac{-1 + \sqrt{-3}}{2} \quad \text{and} \quad \omega^2 = \frac{-1 - \sqrt{-3}}{2} \\
&= \omega^9 + \omega^{18} \\
&= (\omega^3)^3 + (\omega^3)^6 = (1)^3 + (1)^6 && \because \omega^3 = 1 \\
&= 1 + 1 = 2
\end{aligned}$$

(vii)  $\omega^{37} + \omega^{38} - 5$

*Solution:*

$$\begin{aligned}
& \omega^{37} + \omega^{38} - 5 \\
&= \omega^{37} + \omega^{38} - 5 \\
&= \omega^{36} \cdot \omega + \omega^{36} \cdot \omega^2 - 5 \\
&= (\omega^3)^{12} \cdot \omega + (\omega^3)^{12} \cdot \omega^2 - 5 \\
&= (1)^{12} \cdot \omega + (1)^{12} \cdot \omega^2 - 5 \quad \because \omega^3 = 1 \\
&= \omega + \omega^2 - 5 \\
&= -1 - 5 \quad \because \omega + \omega^2 = -1 \\
&= -6
\end{aligned}$$

**(viii)  $\omega^{-13} + \omega^{-17}$**

**Solution:**

$$\begin{aligned}
& \omega^{-13} + \omega^{-17} \\
&= \omega^{-13} + \omega^{-17} \\
&= \omega^{-12-1} + \omega^{-15-2} \\
&= \omega^{12} \cdot \omega^{-1} + \omega^{-15} \cdot \omega^{-2} \\
&= (\omega^3)^{-4} \cdot \omega^{-1} + (\omega^3)^{-5} \cdot \omega^{-2} \\
&= (1)^{-4} \cdot \omega^{-1} + (1)^{-5} \cdot \omega^{-2} \quad \because \omega^3 = 1 \\
&= \omega^{-1} + \omega^{-2} \\
&= \frac{1}{\omega} + \frac{1}{\omega^2} \\
&= \frac{\omega^2 + \omega}{\omega^3} \quad \because \omega^2 + \omega = -1 \text{ and } \omega^3 = 1 \\
&= \frac{-1}{1} \\
&= -1
\end{aligned}$$

**3. Prove that  $x^3 + y^3 = (x + y)(x + \omega y)(x + \omega^2 y)$ .**

**Solution:**

$$\begin{aligned}
x^3 + y^3 &= (x + y)(x + \omega y)(x + \omega^2 y) \\
\text{R.H.S.} &= (x + y)(x + \omega y)(x + \omega^2 y) \\
&= (x + y)[x(x + \omega^2 y) + \omega y(\omega + \omega^2 y)] \\
&= (x + y)[x^2 + \omega^2 xy + \omega xy + \omega^3 y^2] \\
&= (x + y)[x^2 + (\omega^2 + \omega)xy + (1)y^2] \quad \because \omega^3 = 1
\end{aligned}$$

$$= (x + y)[x^2 + (-1)xy + y^2] \quad \because \omega^2 + \omega = -1$$

$$= (x + y)(x^2 - xy + y^2)$$

= L.H.S.

Hence Proved

**4. Prove that  $x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x + \omega y + \omega^2 z)(x + \omega^2 y + \omega z)$ .**

**Solution:**

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x + \omega y + \omega^2 z)(x + \omega^2 y + \omega z).$$

$$\begin{aligned} \text{R.H.S.} &= (x + y + z)(x + \omega y + \omega^2 z)(x + \omega^2 y + \omega z) \\ &= (x + y + z)[x(x + \omega^2 y + \omega z) + \omega y(x + \omega^2 y + \omega z) + \omega^2 z(x + \omega^2 y + \omega z)] \\ &= (x + y + z)[x^2 + \omega^2 xy + \omega xz + \omega xy + \omega^3 y^2 + \omega^2 yz + \omega^2 xz + \omega^4 z + \omega^3 z^2] \\ &= (x + y + z)[x^2 + \omega^2 xy + \omega xy + \omega^2 yz + \omega yz + \omega^2 xz + \omega xz + (1)y^2 + (1)z^2] \\ &= (x + y + z)[x^2 + (\omega^2 + \omega)xy + (\omega^2 + \omega)yz + (\omega^2 + \omega)xz + y^2 + z^2] \quad \because \omega^2 = 1 \\ &= (x + y + z)[x^2 + (-1)xy + (-1)yz + (-1)xz + y^2 + z^2] \quad \because \omega^2 + \omega = -1 \\ &= (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx) \\ &= x^3 + y^3 + z^3 - 3xyz \\ &= \text{L.H.S.} \end{aligned}$$

Hence proved.

**5. Prove that  $(1 + \omega)(1 + \omega^2)(1 + \omega^4)(1 + \omega^8) \dots 2n \text{ factors} = 1$ .**

**Solution:**

$$\begin{aligned} \text{L.H.S.} &= (1 + \omega)(1 + \omega^2)(1 + \omega^4)(1 + \omega^8) \dots 2n \text{ factors} \\ &= (1 + \omega)(1 + \omega^2)(1 + \omega^4)(1 + \omega^8) \dots 2n \text{ factors} \\ &= (1 + \omega)(1 + \omega^2)(1 + \omega^3 \cdot \omega)(1 + \omega^2 \cdot \omega^6) \dots 2n \text{ factors} \\ &= (1 + \omega)(1 + \omega^2)(1 + \omega^3 \cdot \omega)(1 + \omega^2 \cdot (\omega^3)^2) \dots 2n \text{ factors} \\ &= (1 + \omega)(1 + \omega^2)(1 + (1)\omega)(1 + \omega^2(1)^2) \dots 2n \text{ factors} \quad \because \omega^3 = 1 \\ &= (1 + \omega)(1 + \omega^2)(1 + \omega)(1 + \omega^2) \dots 2n \text{ factors} \\ &= (-\omega)^2(-\omega)(-\omega^2)(-\omega) \dots 2n \text{ factors} \quad \because 1 + \omega = -\omega^2 \\ &= [(-\omega)^2(-\omega)][(-\omega^2)(-\omega)] \dots n \text{ factors} \quad \because 1 + \omega^2 = -\omega \\ &= [\omega^3][\omega^3] \dots n \text{ factors} \end{aligned}$$

$$= (1)(1) \dots \text{n factors}$$

$$\therefore \omega^3 = 1$$

$$= (1)^n$$

$$= 1$$

= R.H.S.

Hence proved.

### Roots and co-efficient of a quadratic equation:

We know that  $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$  and  $\frac{-b - \sqrt{b^2 - 4ac}}{2a}$  are roots of the equation

$ax^2 + bx + c = 0$  where  $a, b$  are coefficients of  $x^2$  and  $x$  respectively. While  $c$  is the constant term.

### Relation between roots and co-efficient of a quadratic equation:

$$\text{If } \alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \quad \beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a},$$

then we can find the sum and the product of the roots as follows.

$$\text{Sum of the roots} = \alpha + \beta$$

$$\begin{aligned} &= \frac{-b + \sqrt{b^2 - 4ac}}{2a} + \frac{-b - \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-b + \sqrt{b^2 - 4ac} - b - \sqrt{b^2 - 4ac}}{2a} = \frac{-2b}{2a} = -\frac{b}{a} \end{aligned}$$

$$\text{Product of the roots} = \alpha \beta$$

$$= \left( \frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \left( \frac{-b - \sqrt{b^2 - 4ac}}{2a} \right)$$

$$= \frac{(-b)^2 - (\sqrt{b^2 - 4ac})^2}{4a^2} = \frac{b^2 - (b^2 - 4ac)}{4a^2}$$

$$= \frac{b^2 - b^2 + 4ac}{4a^2} = \frac{4ac}{4a^2} = \frac{c}{a}.$$

If we denote the sum of roots and product of roots by  $S$  and  $P$  respectively, then

$$S = -\frac{b}{a} = -\frac{\text{Co-efficient of } x}{\text{Co-efficient of } x^2} \quad \text{and} \quad P = -\frac{c}{a} = -\frac{\text{Constant term}}{\text{Co-efficient of } x^2}$$

## SOLVED EXERCISE 2.3

- Without solving, find the sum and the product of the following quadratic equations.

$$(i) \quad x^2 - 5x + 3 = 0$$

*Solution:*