

$$\text{Let } \frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$$

$$\Rightarrow \frac{a}{b} = k \quad \text{and} \quad \frac{c}{d} = k \quad \text{and} \quad \frac{e}{f} = k$$

$$a = bk \quad c = dk \quad e = fk$$

$$\begin{aligned} \text{L.H.S.} &= \frac{ac}{bd} + \frac{ce}{df} + \frac{ea}{fb} \\ &= \frac{(bk)(dk)}{bd} + \frac{(dk)(fk)}{df} + \frac{(fk)(bk)}{bf} \\ &= \frac{bdk^2}{bk} + \frac{dfk^2}{df} + \frac{bfk^2}{fb} \\ &= k^2 + k^2 + k^2 \\ &= 3k^2 \quad \text{--- (i)} \end{aligned}$$

$$\begin{aligned} \text{L.H.S.} &= \frac{a^2}{b^2} + \frac{c^2}{d^2} + \frac{e^2}{f^2} \\ &= \frac{b^2k^2}{b^2} + \frac{d^2k^2}{d^2} + \frac{f^2k^2}{f^2} \\ &= k^2 + k^2 + k^2 \\ &= 3k^2 \quad \text{--- (ii)} \end{aligned}$$

From (i) and (ii), we have

$$\text{L.H.S} = \text{R.H.S}$$

$$\text{Hence } \frac{ac}{bd} + \frac{ce}{df} + \frac{ca}{fb} = \frac{a^2}{b^2} + \frac{c^2}{d^2} + \frac{e^2}{f^2}$$

## SOLVED EXERCISE 3.7

1. The surface area  $A$  of a cube varies directly as the square of the length  $l$  of an edge and  $A = 27$  square units when  $l = 3$  units. Find (i)  $A$  when  $l = 4$  units (ii)  $l$  when  $A = 12$  sq. units.

*Solution:*

$$\begin{aligned} \text{Given that } & A \propto l^2 \\ \Rightarrow & A = kl^2 \quad \text{--- (i)} \\ \text{Put } A = 27 \text{ and } l = 3 \text{ in eq. (i), we get} & \\ & 27 = k(3)^2 \\ & 27 = 9k \\ \text{or } & 9k = 27 \\ & k = \frac{27}{9} = 3 \end{aligned}$$

Put  $k = 3$  in eq. (i), we get

$$A = 3l^2 \text{ _____ (ii)}$$

(i) Put  $l = 4$  in eq. (ii), we get

$$\begin{aligned} A &= 3(4)^2 \\ &= 3(16) = \text{eq. units} \end{aligned}$$

(ii) Put  $A = 12$  in eq. (ii), we get

$$\begin{aligned} 12 &= 3l^2 \\ \text{or } 3l^2 &= 12 \\ \Rightarrow l^2 &= 4 \\ l &= 2 \end{aligned}$$

**2. The surface area  $S$  of the sphere varies directly as the square of radius  $r$ , and  $S = 16\pi$  when  $r = 2$ . Find  $r$  when  $S = 36\pi$ .**

**Solution:**

Given that  $S \propto r^2$

$$\Rightarrow S = kr^2 \text{ _____ (i)}$$

Put  $S = 16\pi$  and  $r = 2$  in eq. (i), we get

$$16\pi = k(2)^2$$

Put  $K = 4\pi$  in eq. (i), we get

$$S = 4\pi r^2 \text{ _____ (ii)}$$

Put  $S = 36\pi$  in eq. (ii), we get

$$36\pi = 4\pi r^2$$

$$\text{or } 4\pi r^2 = 36\pi$$

$$r^2 = \frac{36\pi}{4\pi}$$

$$r^2 = 9$$

$$r = 3$$

**3. In Hook's law the force  $F$  applied to stretch a spring varies directly as, the amount of elongation  $S$  and  $F = 3276$  when  $S = 1.6$  in. Find (i)  $S$  when  $F = 50$  lb (ii)  $F$  when  $S = 0.8$  in.**

**Solution:**

Given that  $F \propto S$

$$\Rightarrow F = KS \text{ _____ (i)}$$

Put  $F = 32$  and  $S = 1.6$  in eq. (i), we get

$$32 = k(1.6)$$

$$\text{or } 1.6k = 32$$

$$k = \frac{32}{1.6} = 20$$

Put  $k = 20$  in eq. (i), we get

$$F = 20S \text{ _____ (ii)}$$

(i) Put  $F = 5$  in eq. (ii), we get

$$S = 20 S$$

$$\Rightarrow S = 2.5$$

(ii) Put  $S = 0.8$  in eq. (i), we get

$$F = 20 (0.8) = 16$$

4. The intensity  $I$  of light from a given source varies inversely as the square of the distance  $d$  from it. If the intensity is 20 candlepower at a distance of 12ft. from the source, find the intensity at a point 8ft. from the source.

*Solution:*

Given that  $I \propto \frac{1}{d^2}$

$$\Rightarrow I = \frac{k}{d^2} \text{ _____ (i)}$$

Put  $I = 20$  and  $d = 12$  in eq. (i), we get

$$20 = \frac{k}{(12)^2}$$

$$20 = \frac{k}{144}$$

$$K = 20 \times 144 = 2880$$

Put  $k = 2880$  in eq. (i), we get

$$I = \frac{2880}{d^2} \text{ _____ (ii)}$$

Put  $d = 8$  in eq. (ii), we get

$$I = \frac{2880}{(8)^2} = 45$$

5. The pressure  $P$  in a body of fluid varies directly as the depth  $d$ . If the pressure exerted on the bottom of a tank by a column of fluid 5ft, high is 2.25 lb/sq. in, how deep must the fluid be to exert a pressure of 9 lb/sq. in?

*Solution:*

Given that  $P \propto d$

$$\Rightarrow P = Kd \text{ _____ (i)}$$

Put  $P = 5$  and  $d = 2.25$  in eq. (i), we get

$$5 = K (2.25)$$

$$2.25K = 5$$

$$\Rightarrow K = \frac{5}{2.25} = \frac{20}{9}$$

Put  $k = \frac{20}{9}$  in eq. (i), we get

$$P = \frac{20}{9} d \text{ (ii)}$$

Put  $d = 9$  in eq. (ii), we get

$$P = \frac{20}{9} (9)$$

$$P = 20 \text{ ft}$$

6. Labour costs  $c$  varies jointly as the number of workers  $n$  and the average number of days  $d$ , if the cost of 800 workers for 13 days is Rs. 286000, then find the labour cost of 600 workers for 18 days.

*Solution:*

Given that  $C \propto nd$

$$\Rightarrow C = Knd \text{ (i)}$$

Put  $n = 800$  and  $d = 13$  and  $C = 286000$  in eq. (i), we get

$$286000 = K(800)(13)$$

$$10400K = 286000$$

$$K = \frac{286000}{10400} = \frac{55}{2}$$

Put  $k = \frac{55}{2}$  in eq. (i), we get

$$C = \frac{55}{2} C \text{ (ii)}$$

Put  $n = 600$  and  $d = 18$  in eq. (ii), we get

$$C = \frac{55}{2} \times 600 \times 18$$

$$C = \text{Rs. } 297000$$

7. The supporting load  $c$  of a pillar varies as the fourth power of its diameter  $d$  and inversely as the square of its length  $l$ , A pillar of diameter 6 inch and of height 30 feet will support a load of 63 tons. How high a 4 inch pillar must be to support a load of 28 tons?

*Solution:*

Given that  $C \propto \frac{d^4}{l^2}$

$$\Rightarrow C = k \frac{d^4}{l^2} \text{ (i)}$$

Put  $d = 6$   $l = 30$  and  $C = 63$  in eq. (i), we get

$$63 = K \frac{(6)^4}{(30)^2}$$

$$63 = \frac{1296}{900} K$$

$$\text{or } K = 63 \times \frac{900}{1296}$$

$$K = \frac{175}{4}$$

Put  $k = \frac{175}{4}$  in eq. (i), we get

$$C = \frac{175d^4}{4l^2} \text{ (ii)}$$

Put  $d = 4$  and  $C = 28$  in eq. (ii), we get

$$28 = \frac{175(4)^4}{4l^2}$$

$$l^2 = \frac{175 \times 256}{4 \times 28} = 400$$

$$\Rightarrow l = 20 \text{ ft}$$

8. The time  $T$  required for an elevator to lift a weight varies jointly as the weight  $w$  and the lifting depth ovaries inversely as the power  $p$  of the motor. If 25 sec. are required for a 4-hp motor to lift 500 Ib through 40 ft, what power is required to lift 800 Ib, through 12G ft in 40 sec.?

**Solution:**

Given that  $T \propto Wd$

Also given that  $T \propto \frac{1}{P}$

In joint variation, we can write

$$T \propto \frac{Wd}{P}$$

$$\Rightarrow T = k \frac{Wd}{P} \text{ (i)}$$

Put  $T = 25$ ,  $P = 4$ ,  $W = 500$  and  $d = 40$  in eq. (i), we get

$$25 = \frac{k \times 500 \times 40}{4}$$

$$K = \frac{25 \times 4}{500 \times 40}$$

$$K = \frac{1}{20}$$

Put  $k = \frac{1}{20}$  in eq. (i), we get

$$T = \frac{Wd}{200P} \text{ (ii)}$$

Put  $W = 800$ ,  $d = 120$  and  $T = 40$  in eq. (ii), we get

$$40 = \frac{800 \times 120}{200P}$$

$$P = \frac{800 \times 120}{200 \times 40}$$

$$P = 12 \text{ hp}$$

9. The kinetic energy (K.E.) of a body varies jointly as the mass "m" of the body and the square of its velocity "v". If the kinetic energy is 4320 ft/lb when the mass is 45 lb and the velocity is 24 ft/sec. Determine the kinetic energy of a 3000 lb automobile travelling 44 ft/sec.

Given that  $K.E \propto MV^2$

$$\Rightarrow K.E = KmV^2 \text{ (i)}$$

Put  $K.E = 4320$ ,  $m = 45$  and  $V = 24$  in eq. (i), we get

$$4320 = k(45)(24)^2$$

$$K = \frac{4320}{45 \times 576}$$

$$K = \frac{1}{6}$$

Put  $K = \frac{1}{6}$  in eq. (i), we get

$$K.E = \frac{1}{6} mV^2 \text{ (ii)}$$

Put  $m = 3000$  and  $V = 44$  in eq. (ii), we get

$$\begin{aligned} K.E &= \frac{1}{6} (3000) (44)^2 \\ &= 968000 \end{aligned}$$

## SOLVED MISCELLANEOUS EXERCISE - 3

### 1. Multiple Choice Questions

Four possible answers are given for the following questions. Tick (✓) the correct answer.

(i) In a ratio  $a : b$ ,  $a$  is called

- (a) relation                      (b) antecedent                      (c) consequent                      (d) None of these

(ii) In a ratio  $x : y$ ,  $y$  is called

- (a) relation                      (b) antecedent                      (c) consequent                      (d) None of these

(iii) In a proportion  $a : b :: c : d$ ,  $a$  and  $d$  are called,

- (a) means    (b) extremes  
(c) third proportional                              (d) None of these